

# The Roles of Water Treatment Subsidies and Community Health Workers in Improving Child Health and Chlorine Usage in Malawi

**Sector(s):** Health

**Fieldwork:** Partners In Health (PIH)

**Location:** Southern Malawi

**Sample:** 2313 households from Neno and Mwanza Districts and 427 Community Health Workers (CHWs) from Neno District

**Target group:** Children Extension agents Families and households

**Outcome of interest:** Diarrhea Health outcomes Take-up of program/social service/healthy behavior

**Intervention type:** Subsidies Community health workers

**AEA RCT registration number:** AEARCTR-0002893

**Research Papers:** Expanding Access to Clean Water for the Rural Poor: Experimental Evidence from ...

**Partner organization(s):** Stichting Dioraphte, Stanford Center for Innovation in Global Health, Partners In Health (PIH)

Contaminated water is a carrier of diarrheal disease, the second-leading cause of childhood mortality. While point-of-use treatments like chlorine can be effective at disinfecting drinking water, chlorine usage remains low in sub-Saharan Africa. Researchers conducted a randomized evaluation in Southern Malawi to test the impact of monthly coupons for water treatment solution and different community health worker (CHW) delivery methods on chlorine usage and child health outcomes over an 18-month period. They find that the coupon program had stronger impacts on both outcomes and was more cost-effective than having CHWs distribute free chlorine to households during routine monthly visits.

## Policy issue

Globally, an estimated 1.9 billion people lack access to clean water. In places that cannot afford to provide reliable piped water to rural households, point-of-use water treatment, such as chlorine, can reduce reported child diarrhea by 29 percent and disinfects drinking water against most bacteria. Despite its efficacy, the current usage rate of chlorine among rural households in sub-Saharan Africa is low and finding ways to increase uptake is an ongoing challenge.

One approach to increasing the use of water purification treatments is the free distribution of chlorine through health centers, coupon programs, or Community Health Workers (CHWs). The latter are typically members of the community working either as volunteers or for pay, visiting households and providing basic diagnostic services in association with the local health care system.

While research has demonstrated that targeting via coupons improves the cost-effectiveness of free distribution, this model falls short of achieving widespread coverage of safe water. As numerous countries scale up the use of CHWs, the effectiveness of CHWs in encouraging households to adopt specific health practices remains an open question. Could the CHW model enhance the effectiveness of free chlorine distribution schemes (be it home delivery or coupon programs)?

## Context of the evaluation

Water-borne diseases, such as diarrhea, are a leading cause of death for children under five in Southern Malawi, which may stem from the fact that very few households have access to piped water. Households that participated in the evaluation had mothers

aged 30 on average and just over five members in the family. Only households with a child under the age of six were considered eligible for the study. Sixty-five percent of households used a borehole as their primary source of drinking water and 30 percent used an unprotected source such as a spring.

The evaluation took place in two neighboring districts in Southern Malawi that are relatively low-income and rural: Neno and Mwanza. Researchers collaborated with global health NGO Partners in Health (PIH) to implement the evaluation using their well-established network of CHWs, which operates only in Neno. The CHWs support the government's Health Surveillance Assistants in the community and are each assigned 20 to 30 households to monitor.

The objective of the study was to understand how best to increase usage of WaterGuard, a dilute chlorine solution. One bottle of WaterGuard, if used correctly, would be equivalent to one month's supply of treated water for a family of five. WaterGuard is recognized throughout the study area, although there is widespread concern that it makes the water taste bad. Despite recognition of the product, a survey prior to the program reported that only 4.6 percent of the households in the study area had treated their water with chlorine prior to the intervention.



A water point in Malawi.

Photo credit: Aaron Wolf

## **Details of the intervention**

Researchers partnered with PIH to conduct a randomized evaluation to test the impact of monthly coupons and different CHW delivery methods on chlorine usage and child health outcomes in Southern Malawi. Researchers first randomly assigned 427 CHWs operating in Neno into one of four groups:

1. *WASH Training*: CHWs were trained on how to promote safe drinking water practices, with an emphasis on chlorine use. CHWs in this group were instructed to incorporate chlorine promotion into their monthly household visits.
2. *WASH Training + Home Delivery*: In addition to WASH training, CHWs in this group were instructed to deliver free chlorine directly to households in their catchment area on a monthly basis.
3. *Home Delivery + Rationing*: Eight months after the start of the home delivery treatment, two-thirds of the CHWs from the home delivery group were given enough WaterGuard for only a subset of their assigned households. Rationing was introduced to help reduce wastage and stretch the free delivery program over more months, as data from household surveys suggested that only a minority of households used chlorine. CHWs were explicitly asked to apply their local knowledge and discretion to target households most likely to utilize the free chlorine.
4. *Comparison Group*: CHWs in this group received no special instruction and were expected to continue their regular role in assisting government Health Surveillance Assistants in Neno.

Out of the 1,417 households in Neno whose CHWs had not been assigned to the home delivery group, 709 were randomly assigned to the coupon intervention while the remaining did not receive coupons. In Mwanza, where there was no CHW presence, 440 households were sampled and half were randomly selected to receive coupons. Households in Mwanza and Neno that received coupons could redeem it for a free bottle of WaterGuard at a local shop every month for 18 months. Follow-up surveys on a rolling basis were conducted to trace household chlorine take-up and usage over time. Researchers also collected data on child health outcomes including diarrhea, fever, and vomiting.

## Results and policy lessons

Both the coupon and home delivery programs increased chlorine usage and improved child health outcomes, although the coupon program was more cost-effective than having CHWs distribute free chlorine during routine monthly visits and yielded greater reductions in child illness. Rationing the supply of WaterGuard was more efficient at getting WaterGuard to those who would use it than home delivery to 100 percent of households.

*Coupon take-up*: Although coupon redemption decreased over time from 74 percent in the first month to 40 percent in the last study month, chlorine usage held steady at about 30 percent of coupon households for the duration of the study. This suggests that the drop in take-up in the first few months was among households who had weaker preferences for chlorine use and that self-targeting improved over time. Households with an unprotected source of water redeemed three and a half more coupons on average than those with a protected source. Additionally, households who reported a distaste for chlorinated water prior to the evaluation were less likely to redeem a coupon.

*Chlorine usage*: The coupon intervention across all groups in the two districts increased the likelihood of a positive chlorine test by 26.5 percentage points (a 663 percent increase) while the home delivery intervention increased it by 19.4 percentage points (a 485 percent increase) from a base of 4 percent in the comparison households. Chlorine usage remained at the same level upon introduction of rationing in the home delivery intervention. In Neno, where there was CHW presence, direct delivery by CHWs did not yield higher usage rates than the coupon program, and WASH training by CHWs did not influence overall coupon redemption or chlorine usage in the coupon group.

For households with an unprotected water source prior to the evaluation, the coupon program had a larger impact on chlorine use, with a 36 percentage point increase compared to a 22 percentage point increase in households with a protected water source. In contrast, the effect of the home delivery intervention on chlorine use was not greater for households with unprotected water sources. CHWs visited these households less regularly, likely because they may be located further away.

*Child health*: The likelihood of a child under ten experiencing diarrhea, vomiting, and fever in the past month decreased by 7.6 percentage points (a 18 percent decrease) in the coupon group. The reduction in child illnesses was larger in Mwanza, where child

health was substantially worse on all categories prior to the evaluation. While there was no significant impact of the home delivery interventions on child health outcomes overall, there is evidence of varying effects by season.

*CHW household targeting:* Rationing the supply of WaterGuard was more efficient at getting WaterGuard to those who would use it than home delivery to 100 percent of households. By the last month of the evaluation, the share of households that received a WaterGuard delivery and the share that used it nearly converged in the rationing group, whereas a large gap remained in the 100 percent coverage arm. This suggests that CHWs had private information about household preferences for WaterGuard, which helped them effectively deliver bottles to households that would use them.

*Cost-effectiveness:* Coupons were nearly two times more cost-effective in increasing chlorine usage and four times more cost-effective in averting illness than home delivery with 100 percent coverage. Coupons cost US\$2.69 in 2018 to avert one child illness while home delivery with 100 percent coverage cost US\$10.62 per illness averted.

Together, these results suggest that absent a monitoring or incentive system, targeting through CHWs is less effective than self-targeting through coupon redemption. For low-income families with contaminated water sources, reliable access to water treatment solutions at central locations could be an effective way to target subsidies. This was more cost-effective than relying on a network of CHWs and substantially reduced the incidence of water-borne diseases in low-density, unconnected rural areas, with no decrease in impact over an 18-month period.