

Belief Formation of Risks of Arsenic in Bangladesh

Researchers:

Lori Bennear

Alexander van Geen

Alexander Pfaff

H. B. Soumya

Alessandro Tarozzi

Sector(s): Environment & Energy, Health

Location: Arai hazar Upazila, Bangladesh

Sample: 43 villages

Target group: Rural population

Outcome of interest: Take-up of program/social service/healthy behavior Water access Water quality

Intervention type: Information Preventive health Water, sanitation, and hygiene

AEA RCT registration number: AEARCTR-0001052

Research Papers: Impact of a Randomized Controlled Trial in Arsenic Risk Communication on Househ...

Partner organization(s): Duke University

Policymakers have increasingly seen information disclosure and "labeling campaigns" as a low-cost way to help people avoid environmental exposures, especially in resource-constrained developing countries, yet, there is little evidence on how to present environmental health information in ways that best motivate health-improving behaviors. Researchers compared the effects of two types of arsenic risk information on households' choice of their primary drinking water source. Providing households with more nuanced risk information about arsenic levels in drinking water had no effect on whether they switched to use a safe well.

Policy issue

Environmental exposures like contaminated groundwater have led to widespread health problems in many countries. Policymakers have increasingly seen information disclosure and "labeling campaigns" as a low-cost way to help people avoid environmental exposures, especially in resource-constrained developing countries. If contaminated sites are clearly labeled "unsafe," and people choose to avoid them, information and labeling campaigns could potentially improve their health. Yet, there is little evidence on how to present environmental health information in ways that best motivate health-improving behaviors. Are campaigns that provide more complete, yet complex information about environmental risks more effective than those that give simple safety thresholds?

Context of the evaluation

The natural occurrence of arsenic in groundwater is an important public health concern in Bangladesh, where it is has been estimated that between 35 and 77 million people are at risk of drinking water containing unsafe levels of arsenic, primarily from

shallow tube wells. Exposure to elevated levels of arsenic in water is associated with cancer, developmental effects, cardiovascular disease, skin lesions, deficits in childhood cognitive and motor function, and mortality. No known level of arsenic exposure is completely safe. The most common option to avoid arsenic poisoning in Bangladesh is switching to an uncontaminated well, followed by the construction of deep tube wells. Other options such as piped water systems, rainwater collection, and arsenic filters are used by a very small portion of the population due to their high cost.

The magnitude of the arsenic problem in Bangladesh became evident after almost five million wells were tested by the Bangladesh Arsenic Mitigation and Water Supply Program (BAMWSP) beginning in 1998. The results of the BAMSWP arsenic tests were used to label the wells using a binary system. Wells with arsenic levels in excess of the Bangladesh health standard (50 micrograms per liter) were labeled "unsafe" and painted red, while those with arsenic levels below 50 micrograms per liter were labeled "safe" and painted green. While it has significantly increased well-switching throughout Bangladesh, the binary system may prevent people from making the healthy choice to switch from a highly-contaminated well to a less highly-contaminated well if there is no safe well nearby. Research has shown that health benefits can result from any decrease in arsenic exposure, yet the binary system only encourages switching from a red to a green well.



Collecting drinking water in Bangladesh

Libby Abbott

Details of the intervention

Researchers compared the effects of two types of arsenic risk information on households' choice of their primary drinking water source. In early 2008, wells in 43 villages in Araihasar sub-district were tagged with laser-engraved aluminum plates displaying a pictorial representation of whether the well water was suitable for drinking. The plates show a hand holding a drinking cup that is

covered by a large cross when the arsenic level is above 50 micrograms per liter. It also indicates the year when the water sample was collected and the specific arsenic concentration detected in the water tested from that well. Villages were randomly assigned to receive one of two additional information treatments.

*Status-quo villages:*Received information similar to the information delivered by BAMWSP. Survey staff told households in these villages that, "The federal government says drinking water with more than 50 micrograms arsenic per liter is not safe. When possible you should seek to fetch drinking water from a well that is labeled safe."

*Emphasis villages:*Received additional information that if given a choice between two or more wells, the well with the lowest arsenic level should be chosen, even if they are all marked red or all marked green. Surveyors told households in these villages that, "if you have a choice between a well with 200 micrograms arsenic per liter and a well with 100 micrograms arsenic per liter, drinking water from the well with 100 is better for you between a well with 40 micrograms arsenic per liter and a well with 10 micrograms, drinking water from the well with 10 is better for you. When possible you should seek to fetch drinking water from the well with the lowest arsenic level."

A follow-up survey was conducted in November 2008, between seven and nine months after the information was first delivered.

Results and policy lessons

Providing households with more nuanced risk information about arsenic levels in drinking water had no effect on whether they switched from a contaminated well to a safer one. This result suggests that the additional information treatment did not improve the effectiveness of the status-quo information campaign, and that a one-time oral communication about the health benefits of switching to a less contaminated well will not be sufficient to increase switching from highly-contaminated wells to less contaminated wells.

One primary policy implication of this finding is that messages conveying richer information on arsenic risks, while inexpensive and easy to scale, are unlikely to be successful in reducing exposure relative to the status-quo policy of providing a simple indicator that a well is safe or unsafe. However, a more sophisticated presentation of the risk spectrum on the aluminum plates installed on wells, showing gradients that represent the level of arsenic in the well relative to others in the village, may have greater potential to induce well-switching.

Ahmed, Kazi Matin, Lori Bennear, Alexander Pfaff, H. B. Soumya, Alessandro Tarozzi, and Alexander van Geen. 2013. "Impact of a randomized controlled trial in arsenic risk communication on household water-source choices in Bangladesh." *Journal of Environmental Economics and Management* 65(2): 225-240.